

OFFICIAL

comparing the agent-biased binding with a corresponding unbiased binding of the sensor to the first receptor;

wherein a difference between the biased and unbiased bindings indicates that the agent modulates a binding function of the first receptor.

33. (New) A method according to claim 32, wherein the measuring step comprises detecting the first receptor of immobilized first receptor-sensor complexes.

34. (New) A method according to claim 32, wherein the measuring step, the first receptor is immobilized through the sensor.

35. (New) A method according to claim 32, wherein the sensor comprises a label and wherein the measuring step, the first receptor is immobilized through the sensor and the sensor is immobilized through the label.

36. (New) A method according to claim 32, wherein the sensor comprises a label and wherein the measuring step, the first receptor is immobilized through the sensor, and the sensor is immobilized through the label by a second receptor.

37. (New) A method according to claim 32, wherein the sensor comprises a label and wherein the measuring step, the first receptor is immobilized through the sensor, and the sensor is immobilized through the label by a second receptor and wherein the measuring step comprises detecting the immobilized first receptor.

38. (New) A method according to claim 32, wherein the sensor comprises a label and wherein the measuring step, the first receptor is immobilized through the sensor, and the sensor is immobilized through the label by a second receptor and wherein the measuring step comprises detecting the immobilized first receptor with a third receptor.

39. (New) A method according to claim 32, wherein the sensor comprises an epitope label, wherein the measuring step, the first receptor is immobilized through the sensor and the sensor is

OFFICIAL

immobilized through the label by a second receptor comprising an immobilized epitope label-specific antibody moiety.

40. (New) A method according to claim 32, wherein the sensor comprises a biotin label and wherein the measuring step, the first receptor is immobilized through the sensor and the sensor is immobilized through the label by a second receptor comprising an immobilized avidin moiety.

C1
Cent'd
41. (New) A method according to claim 32, wherein the measuring step, the sensor is immobilized through the first receptor.

42. (New) A method according to claim 32, wherein the measuring step, the sensor is immobilized through the first receptor and the first receptor is immobilized through a second receptor.

43. (New) A method according to claim 32, wherein the measuring step, the sensor is immobilized through the first receptor and the first receptor is immobilized through a second receptor and wherein the measuring step comprises detecting the immobilized sensor.

44. (New) A method according to claim 32, wherein the measuring step, the sensor is immobilized through the first receptor and the first receptor is immobilized through a second receptor and wherein the measuring step comprises detecting the immobilized sensor with a third receptor.

45. (New) A method according to claim 32, wherein the measuring step, the sensor is immobilized through the first receptor and the first receptor is immobilized through a second receptor comprising a receptor specific antibody.

46. (New) A method according to claim 32, wherein the first receptor comprises the ligand binding domain of PPAR γ , Cyp7PBP(LRH-1), NURR1, RZR β , ROR α , NOR-1, Rev-ErbA β , Tlx, NGFI-B β , HZF-2 α , COUP-TF α , β , γ , Nur77, LXR α , COR, Rev-ErbA α , HNF4 α , TOR, MB67 α , SHP, FXR, SF-1, LXR β , GCNF, TR2-11 α , β , TR4, ERR α , β and DAX-1

OFFICIAL

47. (New) A method according to claim 32, wherein the agent effects an increase in binding of the sensor to the first receptor.
48. (New) A method according to claim 32, wherein the sensor is at a concentration of less than about 10 nM.
49. (New) A method according to claim 32, wherein the first receptor, peptide and agent are in solution.
50. (New) A method according to claim 32, wherein the peptide comprises a fluorescent label and the measuring step comprises detecting fluorescence polarization of the label.
51. (New) A method according to claim 32, wherein the mixture further comprises a ligand of the first receptor.
52. (New) A method according to claim 32, wherein the peptide is 12 or fewer residues in length.
53. (New) A method according to claim 32, wherein the peptide comprises an amphipathic alpha helix.
54. (New) A method according to claim 32, wherein the peptide comprises a sequence selected from the group consisting of: KLVQLTTT (SEQ ID NO:1), ILHRLLE (SEQ ID NO:2), LLRYLLDK (SEQ ID NO:3), LLRYLLD (SEQ ID NO:4), LRYLLD (SEQ ID NO:5), LLRYLL (SEQ ID NO:6), LRYLL (SEQ ID NO:7), LLRYLLDKD (SEQ ID NO:8), QLLRYLLDKD (SEQ ID NO:9), HQLRYLLDKD (SEQ ID NO:10), PQAQQKSLLQQLLT (SEQ ID NO:11), LLQQLLTE (SEQ ID NO:12), VTLLQLLG (SEQ ID NO:13), ILRKLLQ (SEQ ID NO:14), ILKRLLE (SEQ ID NO:15), ILRLLQ (SEQ ID NO:16) and ILKLLQ (SEQ ID NO:17).
55. (New) A method according to claim 32, wherein the peptide consists of a sequence selected from the group consisting of: KLVQLTTT (SEQ ID NO:1), ILHRLLE (SEQ ID NO:2),

OFFICIAL

LLRYLLDK (SEQ ID NO:3), LLRYLLD (SEQ ID NO:4), LRYLLD (SEQ ID NO:5), LLRYLL (SEQ ID NO:6), LRYLL (SEQ ID NO:7), LLRYLLDKD (SEQ ID NO:8), QLLRYLLDKD (SEQ ID NO:9), HQLRYLLDKD (SEQ ID NO:10), PQAQKSLQQLLT (SEQ ID NO:11), LLQQLLTE (SEQ ID NO:12), VTLLQLLG (SEQ ID NO:13), ILRKLLQE (SEQ ID NO:14), ILKRLLQE (SEQ ID NO:15), ILRRLLQE (SEQ ID NO:16) and ILKKLLQE (SEQ ID NO:17).

56. (New) A method according to claim 32, wherein the label provides for indirect detection of the sensor.

57. (New) A method according to claim 32, wherein the label provides for indirect detection of the sensor, wherein the label is an epitope tag.

58. (New) A method according to claim 32, wherein the label provides for direct detection of the sensor.

59. (New) A method according to claim 32, wherein the label provides for direct detection of the sensor, wherein the label is a luminescent label.

60. (New) A method according to claim 32, wherein the label provides for direct detection of the sensor, wherein the label is a luminescent label, wherein the luminescent label is a fluorescent label.

61. (New) A method according to claim 32, wherein the label provides for direct detection of the sensor, wherein the label is a luminescent label, wherein the luminescent label is a fluorescent label, wherein the fluorescent label is coupled to the N-terminus of the peptide.

REMARKS

New claims 32-45 are similar to canceled claims 17-30; new claims 46-55 are similar to canceled claims 2-5, 7-10 and 16. Support for labels providing for direct and indirect detection (claims 56-61) is found on p.7, lines 12-20). Consistent with the amendments made today in copending Serial No. 08/975,614, the "5-25" limitation of our July 17 amendment has been